



RESEARCH PAPER

Response of supplemental cassava root sievate - cassava leaf meal based diets on carcass and cost benefits of kano brown goats *Capra aegagrus*

Ahaotu E. Obioha¹, Okorie Kennedy^{2*}, Akinfemi Abayomi³, Emeribe E. Okechukwu⁴

¹Department of Animal Production and Health Technology, Imo State Polytechnic Umuagwo, Nigeria; ²Department of Animal Science and Fisheries, Imo State University, P.M.B 2000, Owerri, Nigeria;

³Department of Animal Science, Nasarawa State University, Lafia, Nigeria; ⁴Department of Crop Science and Biotechnology, Imo State University Owerri, Nigeria. Email: oshotex2014@yahoo.ca

ABSTRACT

The experiment was conducted to determine the organ weight characteristics, carcass yield and cost benefits of Kano Brown goats fed cassava root sievate and cassava leaf meal (CRSCLM) based diets as supplement to (*Pennisetum purpureum*) using 36 Kano Brown goats of 8 to 10 months of age. Four diets T₁, T₂, T₃, and T₄, were formulated at the levels of 0%, 20%, 40% and 60% CRSCLM respectively in a completely randomized design. Each animal received a designated treatment diet in the morning for 97 days. Feed offered was based on 3.5% body weight per day; the animals in addition were fed a kg wilted chopped *P. purpureum* later in the day as basal diet to enhance rumination and fine chewing. Results on carcass indices showed significant ($p < 0.05$) response on live weight at slaughter, empty carcass weight, warm carcass weight, dressing percentage, shoulder, leg, loin, end and shank with T₄ having relatively best results. The dressing percentage was numerically (49.59%) best at T₄. On the offal weights, head and full guts were significantly ($p < 0.05$) improved for T₄ and T₁ goats. The organ characteristic proved the safety of using CRSCLM through the significantly ($p < 0.05$) lower organ weights at T₄ goats. Cost per kg feed, feed cost/weight gain and cost/benefit ratio were positively influenced ($p < 0.05$) at T₄ with expected income of 4.79 / 1 invested. It could therefore be concluded that Kano Brown goats fed 60% CRSCLM had the best carcass and organ yields at a reduced feed cost.

Keywords: *Pennisetum purpureum*, Kano brown goats, Cassava leaf meal, Cassava root sievate,

INTRODUCTION

Livestock production is an important tool in the economy of developing countries. Goats particularly play an important role in the livelihood of small scale farmers as a major component of livestock mixed farming systems, which produce meat, milk, skin, fiber and manure to large number of low income earners (Ahaotu and Ayo-Enwerem, 2018). The products are however in short supply due to seasonal fluctuation in feed quality and quantity, thus making feeding an important constraint to improved goat production.

The provision of adequate nutrition could thus be a pathway to enhanced goat production. Grasses which are the most abundant basal feed for goats dry up during the dry season thus resulting to highly lignified feed material with very poor digestibility coefficient, low energy and protein values. The complex factor that affects livestock production is nutrition, as feed resources are limited in quantity and quality (Ahaotu, 2018). The systems of goat production in Nigeria are usually characterized by limitations posed by non-availability of year-round feed resources due to prolonged dry season of northern Nigeria (Aina *et al.*, 2002). This will in turn lead to poor performance of the animals (Katuromunda *et al.*, 2012). This however has encouraged the need for a dry season supplementation to augment for the shortfall in the deficient nutrients so as to meet the nutrient requirements of goats. One of the readily, abundant

and cheaply sourced feedstuff which could be used to formulated a dry season supplement for goats is cassava root sievate and cassava leaf meal due to their high energy, fibre and protein profile.

The supplemental diets must be formulated in such a way not to compromise the nutritional need of the goats, but should be however be at affordable cost. The use of cassava or its by-products in the formulation of supplemental diets for ruminants has been extensively researched. Goats (Oni *et al.*, 2012; Katurumunda *et al.*, 2012; Oni *et al.*, 2017), sheep (Fasae *et al.*, 2015; Fadiyimu *et al.*, 2016) and cattle (Oduguwa *et al.*, 2013) fed with cassava or its by-products showed improved feed intake, nutrient utilization, blood profile, growth rate, carcass and organ characteristics. Cassava root sievate and cassava leaf meal have been identified as an underutilized feed resources for goats, its nutritional quality has necessitated its utilization as goat feed. Cassava leaf meal is a non-conventional feed stuff that is readily available at little or no cost. The root sievate has a high energy and fibre profile which are essential for goats. Cassava and its by products have been however, implicated with anti-nutritional factor majorly cyanide (Morgan and Choct, 2016) which has been reported to affect nutrient availability, blood formation and utilization when not properly processed (Oelker *et al.*, 2009). Thus, the fermentation, drying and further milling employed in this study would perhaps further reduce the anti-nutrients to tolerable level for goats. This experiment was designed therefore to determine the carcass yield, organ weights and economics of production of Kano brown goats fed cassava root sievate-cassava leaf meal based diets.

MATERIALS AND METHODS

The experiment was carried out at the Goatery Unit of Animal Production and Health Technology Department, Imo State Polytechnic Umuagwo, Nigeria. Imo State Polytechnic Umuagwo is located at the site is situated between longitudes 7° 01' 06¹¹E and 7° 03' 00¹¹ and latitudes 5° 28' 00¹¹N and 5° 30' 00¹¹N in the humid tropical West Africa (IMLS, 2016). The climate is marked by two seasons.

Experimental feeds

The cassava root sievate and cassava leaf were sourced and harvested within the Imo State Polytechnic Communities, Umuagwo, Nigeria. The cassava root sievate is a by-product of cassava root processing. It was gotten after the processed cassava roots production are peeled or not, washed clean and soaked in clean water for 3-5 days to ferment so as to reduce the hydrogen cyanide level and also to soften the roots to enable sieving. Thereafter, the soaked cassava roots are sieved, the sievate (waste) collected and sundried for about 5 days to reduce the moisture contents and possible anti-nutrients that were not removed during the fermentation process. The cassava leaves were harvested from the Department of Crop Production research and demonstration cassava farms after root harvesting. The both were coarsely milled using hammer mill to encourage fine chewing. The cassava root sievate meal (CRSM) and cassava leaf meal (CLM) were mixed in the ratio of 3:1 and used in the formulation of the experimental diet. Four diets T₁, T₂, T₃, and T₄, were formulated. The cassava root sievate-cassava leaf meal (CRSCLM) was included at the levels of 0%, 20%, 40% and 60% for T₁, T₂, T₃, and T₄, respectively as presented in Table 1. All feeds and test ingredients were analyzed for proximate compositions using the method of AOAC (2000).

Animal management

Thirty six (36) Kano Brown goats within the range of 8 – 10 months of age and averaging 7.19kg in weight were selected from the Imo State Polytechnic goatery farm for this experiment. The goats were randomly divided into four groups of nine animals each with three goats

constituting a replicate. The groups were randomly assigned the four experimental diets (T₁, T₂, T₃, and T₄) in a completely randomized design (CRD).

The animals were housed individually in a well-ventilated cement floored pens equipped with feeders and drinkers. Each animal received a designated treatment diet in the morning (8am) for 97 days. Feed offered was based on 3.5% body weight per day; the animals in addition were fed 1kg wilted chopped *P. purpureum* later in the day (5pm). Regular access to fresh drinking water was made available.

Table 1. Composition of the experimental diets for Kano Brown Goats

Ingredients	Dietary levels (%)			
	T ₁	T ₂	T ₃	T ₄
CRSCLM	0.00	20.00	40.00	60.00
PKM	48.00	38.00	30.00	20.50
BDG	47.50	37.50	25.50	15.00
Molasses	2.00	2.00	2.00	2.00
Bone meal	1.00	1.00	1.00	1.00
Salt	0.50	0.50	0.50	0.50
Limestone	1.00	1.00	1.00	1.00
Total	100	100	100	100

Table 2. Proximate composition of the experimental diets, cassava root sievate meal, cassava leaf meal and *Pennisetum purpureum*

Nutrients (%)	Treatments						
	T ₁ (0)	T ₂ (20)	T ₃ (40)	T ₄ (60)	CRSM	CLM	PP
Dry matter	89.95	90.40	91.00	91.44	88.60	89.12	30.93
Crude protein	15.36	14.87	13.64	13.00	2.57	17.66	5.34
Crude fibre	18.96	18.96	19.65	20.11	18.96	5.38	12.64
Ash	5.44	5.59	4.91	4.50	1.80	9.87	4.01
Ether Extract	2.15	3.26	3.38	2.90	2.71	3.93	3.17
Nitrogen Free Extract	48.04	42.72	50.14	50.93	68.14	52.28	26.37
Gross Energy (MJ/kg)	3.94	3.89	4.07	4.04	3.79	3.76	2.27
Neutral Detergent Fibre	62.44	42.08	35.75	33.93	25.34	39.90	58.31
Acid Detergent Fibre	58.35	38.30	25.11	23.69	6.68	33.25	28.60
Hemi Cellulose	4.09	3.78	10.64	10.24	8.66	6.65	19.17

CRSM = Cassava root sievate meal; CLM = cassava leaf meal; PP = *Pennisetum purpureum*

Carcass and organ evaluation

After the 97 days feeding trial, one goat from each replicate was starved of feed for 24 hours and weighed prior to slaughter. The goats were cut at the throat, and then slaughtered by severing the heads at its articulation with the atlas bone. After dressing, the carcasses were weighed to determine the warm carcass weight. This represented the weight of the goats after removal of head, skin, thoracic, abdominal and pelvic contents and the limb distal to the joints. The dressing out percentages were calculated by dividing the warm carcass weight by the live weight prior to slaughter and multiplied by 100%. Other carcass components, organ, guts and muscles were weighed as well. The empty weights were determined by subtracting the weights of the gut contents from the live weights at slaughter. The warm carcasses were divided into two with a saw through the spinal column. The left halves were cut into various parts. The leg/thigh was severed from the attachment of the femur to the acetabulum. The loin consisted of the lumber region plus a pair of ribs and the ends (9 spare ribs and the belly) of six abdominal ribs. The shoulder consisted of the scapula, humerus, radius, ulna, carpals while the sets are made up of the breast and the neck. Each of the cut parts was weighed and doubled in each case before being expressed as a percentage of warm carcass weight.

Economics of production

The variable cost of feeding the goats considered as the cost of the feed and all other cost (i.e. labour, capital investment and housing) were the same for all the treatments. The cost of processing the cassava root sievate-cassava leaf meal (CRS-CLM) was included as the feed cost. The cost per 100kg feed (#) was determined by the summation of the cost of each feed ingredient used in the formulation per treatment. The cost per kilogramme of feed (#) was determined by dividing the cost per 100kg feed by 100. Total cost of feed (#) was calculated by multiplying total feed consumed by cost per kg of feed and daily feed cost was calculated by dividing total feed cost by the number of experimental days. Feed cost/weight gain (#) was determined by dividing total cost of feed by total weight gain. Cost benefit ratio was calculated by dividing cost/kg live weight by feed cost/weight gain.

Statistical analysis

The results were analyzed using one-way analysis of variance (ANOVA) to determine the means and standard error. Treatment means were compared using Duncan's new multiple range test (Gordon and Gordon, 2004).

RESULTS

The proximate composition of the experimental diets, CRSM, CLM and *P. purpureum* are presented in Table 2. The dry matter (DM), crude fibre (CF) and gross energy increased with increasing levels of CRSCLM while crude protein (CP), Neutral detergent fibre (NDF) and Acid detergent fibre (ADF) decreased with increasing levels of CRSCLM. Ash, ether extract (EE), Nitrogen free extract (NFE) and hemicellulose failed to follow a specific trend across the treatment groups. Diets had significant ($p < 0.05$) effects on all the carcass parameters (Table 3), except for set that remained significantly ($p > 0.05$) similar across the treatments. T₃ had the lowest slaughter weight which significantly ($p < 0.05$) differed from T₄. The difference in the slaughter weight may be attributed to lower initial weight of T₃ animals or may be due to high final weight observed for T₄ goats. T₃ showed significantly ($p < 0.05$) lower empty weight in comparison to other treatments. The meat cuts are expressed as percentage of warm carcass weight apart from set, were influenced ($p < 0.05$) by the dietary levels.

The fibre fractions (NDF, ADF and hemicellulose) have implication on the digestibility of plants. The NDF, which is a measure of the plants' cell wall contents, is the chemical component of the feed that determines its rate of digestion.

Table 3. Carcass characteristics of kano brown goats fed cassava root sievate-cassava leaf meal based Diets

Parameters	T ₁	T ₂	T ₃	T ₄	SEM
Live weight at slaughter (kg)	13.17 ^a	13.51 ^a	11.90 ^b	14.46 ^a	0.38
Empty carcass weight (kg)	10.57 ^a	9.99 ^a	7.75 ^b	11.35 ^a	0.78
Warm carcass weight (kg)	6.20 ^a	5.67 ^b	4.43 ^b	7.17 ^a	0.63
Dressing percentage (%)	47.08 ^a	41.97 ^b	37.23 ^b	49.59 ^a	3.19
Meat cuts expressed as percentage (%) of warm carcass weight					
Shoulder	9.23 ^b	10.39 ^b	8.58 ^b	13.35 ^a	1.80
Leg	21.13 ^b	34.73 ^a	19.40 ^b	34.40 ^a	22.26
Loin	23.23 ^b	24.66 ^b	22.01 ^b	27.61 ^a	2.51
Sets	22.80	24.70	19.69	20.79	2.54
End	3.98 ^c	5.35 ^b	6.66 ^a	6.84 ^a	0.57
Shank	12.31 ^b	12.45 ^b	12.45 ^b	17.32 ^a	2.21

^{a-c} means in the row with different superscripts are significantly different ($p < 0.05$)

Table 4. Offal characteristics of kano brown goats fed cassava root sievate-cassava leaf meal based diets

Parameters	T ₁	T ₂	T ₃	T ₄	SEM
Head	11.55 ^a	12.97 ^a	8.68 ^b	13.97 ^a	1.78
Full gut	16.85 ^a	13.23 ^b	12.07 ^b	16.60 ^a	2.31
Feet	6.30	6.66	7.76	7.51	0.30
Skin	8.00	7.56	7.16	6.54	0.31
Empty gut	5.39	5.44	6.59	5.68	0.16

^{a-b} means in the row with different superscripts are significantly different ($p < 0.05$)

The DM, CP, ash, NDF, ADF and hemicellulose were higher in the control, but however compared to the treatment groups.

Table 5. Organ characteristics of kano brown goats fed cassava root sievate-cassava leaf meal based diets

Parameters	T ₁	T ₂	T ₃	T ₄	SEM
Spleen	0.985	1.17	1.375	1.14	0.03
Liver	3.505 ^b	3.640 ^b	5.725 ^a	2.070 ^b	0.81
Heart	1.230 ^b	1.170 ^b	1.925 ^a	0.855 ^b	0.07
Kidney	0.915 ^b	1.110 ^b	2.640 ^a	0.855 ^b	0.22
Testes	1.215	0.59	1.445	1.695	0.12
Lungs	1.290 ^a	1.110 ^a	1.805 ^a	0.550 ^b	0.11

^{a-b} means in the row with different superscripts are significantly different ($p < 0.05$)

Table 6. Cost benefit analysis of kano brown goats fed supplemental cassava root sievate - cassava leaf Meal

Parameters	T ₁	T ₂	T ₃	T ₄	SEM
Cost/100kg feed (#)	4975.00 ^a	4325.00 ^a	4120.00 ^b	3854.50 ^b	69.21
Cost/kg of feed (#)	49.75 ^a	43.25 ^b	41.20 ^b	38.55 ^b	0.69
Total feed consumed (kg)	31.54	33.15	32.71	34.27	1.17
Total cost of feed (#)	1569.12 ^a	1433.74 ^b	1347.65 ^c	1321.11 ^c	47.1
Daily feed cost (#)	16.18 ^a	14.78 ^b	13.89 ^c	13.62 ^c	0.46
Total weight gain (kg)	4.76 ^b	4.67 ^b	5.04 ^{ab}	5.27 ^a	0.32
Feed cost/weight gain (#)	329.65 ^a	307.01 ^a	267.39 ^b	250.69 ^b	8.35
Cost/kg live weight (#)	1200.00	1200.00	1200.00	1200.00	0.00
Cost benefit ratio	1:3.64	1:3.91	1:4.49	1:4.79	0.00

^{a-c} means in the row with different superscripts are significantly different ($p < 0.05$)

(#) indicated the Naira; Nigerian currency.

DISCUSSION

The empty weight (7.75-11.35kg) obtained in this study is lower than 13.09-19.72kg reported by Marques *et al.* (2014) for Moxotó growing goats, but somehow comparable with 8.1-9.2kg reported by Grum (2010) for Kano brown goats. The warm carcass weight (4.43-7.17kg) obtained in this study is lower than 11.01kg reported by Rodriguez *et al.* (2014) and higher than 3.40-4.65kg reported by Ifut *et al.* (2011).

The carcass weights are however in agreement with the findings of Ifut *et al.* (2011) who reported significant values in warm carcass weights. The values obtained in this study also fell within the range of 2.9-9.81kg reported by Marichal *et al.* (2003). Ahaotu and Ayo-Enwerem (2008) stated that dressing percentage (DP) is affected by plane of nutrition and other factors such as breed, age and sex. However, the significant ($p < 0.05$) differences observed in this study may be attributed to nutrition, age and slaughter weight of the goats. The dressing percentage obtained in this study however, fell within 38 and 56% baseline reported for goats. This is an indication that the diets were nourishing and supported lean meat deposition across the treatment groups. The effect of nutrition on dressing percentage cannot be over emphasis because changes

in goat diets may improve both the quantity and quality of the goat meat as a final product (Geay *et al.*, 2001). The influence of nutrition, age and slaughter weights has been reported to positively influence dressing percentages in goats.

The meat cut is an important factor in evaluation of meat yield of goat. T₄ diet showed a superior shoulder, leg, loin, and shank weights than other treatment groups. This suggested that T₄ diet influenced the development of these cut parts better than the other treatment groups. The factors that may have attributed to the variations in the shoulder, leg, loin and shank weights include age, slaughter weight and body condition of the goats. The improved meat cut parts at T₄ may be attributed to higher available energy and lower ADF, NDF and lower protein content of the diet. High percentages of crude fibre and roughage with low digestibility could contribute to low dressing percentages which is a function of the meat cuts. Mahgoub *et al.* (2005) noted that increased level of protein supplement resulted to decrease in the dressing percentage (meat cuts) while increased ME levels resulted to increased carcass weight and DP.

The edible offal weights of Kano brown goats fed CRSMCLM based diets are presented in Table 4. T₁, T₂ and T₄ showed significantly ($p < 0.05$) superior head values than T₃ goats. Table 5 summarizes the organ yields of Kano brown goats fed *P. purpureum* supplemented with CRSMCLM based diets. The significant ($p < 0.05$) value observed in this present study for head is in agreement with Ifut *et al.* (2011) for Kano brown goats. The full gut also differed significantly ($p < 0.05$) with T₃ having the lowest value. The result however, is in disagreement with Anbarasu, (2004) who reported that gut fill could constitute 20 -22% of live weight.

There was significant difference ($p < 0.05$) in the organ yields of the treatments evaluated except for spleen weights. It is a common practice in feeding trials to use weight of some internal organs like kidney and liver as indicators of anti-nutritional factors (Ahaotu and Ayo-Enwerem, 2008). Ahaotu (2018) reported that if there is any toxic element in feed samples used in feeding trial, abnormalities will arise because of increased metabolic activities of the organ in an attempt to reduce these toxic elements or anti-nutritional factors to non-toxic forms.

The cost benefit evaluation of Kano brown goats fed CRSCLM in their diets is presented in table 6. The cost of production per 100 kg feed, cost per kg of feed, total cost of feed and daily feed cost differed ($p < 0.05$) significantly across the treatment groups and tended to decrease with increasing levels of CRSCLM inclusion in the diets. As the level of inclusion of CRSCLM in the diets of Kano brown goats increased from 0% (T₁) 0% to 60% (T₄), the overall cost of feed decreased. The feed cost/weight gain showed significant ($p < 0.05$) values decreasing with increasing levels of CRSCLM. The decrease in feed cost maybe attributed to the inclusion of locally non-competitive and very cheap unconventional feed (CRSCLM) in the diets of Kano brown goats. This is in agreement which earlier reports by Katurumunda *et al.* (2012) who reported a reduced feed cost when *Moringa oleifera* leaf meal (locally and cheaply sourced alternative) was included in the diets of Kano brown goats. From the result, it could be deduced that CRSCLM also enhanced weight gain which further resulted to lower feed cost per kg weight gain. A similar result was reported by Katurumunda *et al.* (2012). This further entail that inclusion of some potential locally cheap feed stuffs do not only reduce feed cost, but also increases weight gain. Cost benefit ratio was best for T₄. This was agreement with the results of earlier studies by (Ogundipe *et al.*, 2003) who reported that the need to lower feed cost in order to produce affordable meat and other animal products for the populace cannot be over-emphasized in the face of dwindling standard of living. The result demonstrated the qualitative benefits and financial returns of using CRSCLM diets; with T₄ having the highest ratio and T₁ had the lowest value. This entails an expected benefit of # 4.79 for every # 1 in cost for T₄ diet. This result suggested that the optimum level of inclusion of CRSCLM in the diet of Kano brown goat may not have been attained and perhaps incremental level beyond 60% inclusion may still yield higher cost benefit ratio beyond the value recorded for T₄ in this trial.

The proximate composition of *P. purpureum* in this study is in comparison with the values reported by Onyeonagu and Eze (2013) and Amaefule and Elendu (2010) for the same forage. The cassava leaf meal (CLM) is comparable with the DM, CP, ash and NFE values reported by

Allen (1984) and Akinfala *et al.* (2002). The crude protein content of the CRSM is below the acceptable 7% CP for ruminant performance as recommended by Ahaotu (2018) and 8% suggested by Ahaotu and Ayo-Enwerem, (2008) for ruminal function.

The NDF comprises mainly the cell wall fraction of forages and roughages and includes a complex matrix of lignin, small amounts of protein, and various polysaccharides, Ocheja *et al.*, (2014) noted that the higher the NDF, the lower the plant's digestible energy. The values obtained for the CRSM may imply moderate cell wall content, moderate digestible energy and DM intake. The ADF consist mainly the lignin and cellulose. Hemicellulose has been reported to be more digestible than cellulose (Ihezuo *et al.*, 2013). The reportedly lower values of the fibre fractions are in agreement with the findings of Ajeigbe *et al.* (2011) for the same cassava by product. The high energy value reported for the CRSM is in agreement with Khampa *et al.* (2009) who noted that cassava roots contain high levels of energy and have been used as a source of readily fermentable energy in ruminant rations. The high dry matter value reported is favourably compared with the values of Boonnop *et al.* (2009). The proximate composition of the experimental diets revealed that the crude protein and the energy requirements are within the ranges reported for goats (Ahaotu and Ayo-Enwerem, 2008).

CONCLUSIONS

This study revealed that the inclusion of cassava root sievate – cassava leaf meal in the diets of Kano brown goats resulted to superior carcass and organ weights and lower production cost with high return on investment. It further revealed that the processing methods used in this study reduced the level of anti-nutritional factors (HCN) to a tolerable level by Kano brown goats. It was therefore concluded that Kano brown goats fed 60% cassava root sievate – cassava leaf meal had the best carcass and organ yields at a reduced feed cost.

REFERENCES

- Ahaotu, E.O. 2018. Sheanut vitellaria paradoxa cake as ingredient in the diet of west African dwarf sheep *Capra ovis*. Aceh Journal of Animal Science, 3 (2): 47-54.
- Ahaotu, E.O., C.M. Ayo-Enwerem. 2008. Goat production and management practices. Jeolas Press, Owerri, Nigeria.
- Ajeigbe, H.A., B.B. Singh, A. Musa, S.A. Tarawali. 2011. Effect of feeding crop residues of different cereals and legumes on weight gain of Yankasa rams. International Journal of Livestock Production, 2(2): 17-23.
- AOAC. 2000. Association of Official Analytical Chemists: Official Methods of Analysis. 6th Edition. Washington DC, USA.
- Anbarasu, C., N. Dutta, K. Sharma, M. Rawat. 2004. Response of goats to partial replacements of dietary protein by a leaf meal mixture containing *Leucaena leucocephala*, *Morus alba* and *Tectona grandis*. Small Ruminant Research, 51: 47-56.
- Boonnop, K., M. Wanapat, N. Nontaso, S. Wanapat. 2009. Enriching nutritive value of cassava root by yeast fermentation. Science Agriculture, 66: 629-633.
- Fadiyimu, A., J.A., Alokun, A.N. Fajemisin, G.E. Onibi. 2016. Feed intake, growth performance and carcass characteristics of west african dwarf sheep fed *Moringa oleifera*, *Gliricidia sepium* or cassava fodder as supplements to *Panicum maximum*. Journal of Experimental Agriculture International, 14: 1-10.
- Fasae, O.A., A.O. Amos, A. Owodunni, A.O. Yusuf. 2015. Performance, haematological parameters and faecal egg count of semi-intensively managed west African dwarf sheep

- to varying levels of cassava leaves and peels supplementation. *Pertanika Journal of Tropical Agricultural Science*, 38: 71 – 81.
- Geay, Y., D. Bauchart, J.F. Hocquette, J. Culioli. 2001. Effect of nutritional factors on biochemical, structural and metabolic characteristics of muscles in ruminants, consequences on dietetic value and sensorial qualities of meat. *Reproduction Nutrition Development*, 41: 1-26.
- Gordon, S.P., F.S. Gordon. 2004. *Contemporary statistics: A computer approach*. McGraw – Hill Publishers, U.S.A.
- Grum, G. 2010. Community-based participatory characterization of the short eared somali goat population around dire dawa. MSc thesis, School of Animal and Range Science, School of Graduate Studies Haramaya University, Haranaya.
- Ihezuo, J.P, E.O. Ahaotu, R.E. Uwalaka. 2013. Replacement of rice and maize stalks with *Tephrosia* spp. and *Moringa oleifera* cakes in the diet of small ruminants raised under humid tropics. *International Journal Applied Sciences and Engineering*, 1(2): 66-68.
- Ifut, O.J., U.A. Inyang, I.S. Udosi, M.I. Ekpo. 2011. Carcass yield of west african dwarf goats fed mixed forages and brewers' spent grain. *Nigerian Journal of Agriculture, Food and Environment*, 7(2):77-83.
- IMLS. 2016. Imo State of Nigeria Ministry of lands and survey – Detailed Atlas of the State.
- Katuromunda, S., E.N. Sabiiti, A. Mateete-Bekunda. 2012. Effect of legume foliage supplementary feeding to dairy cattle offered *Pennisetum purpureum* basal diet on feed intake and manure quality. *Uganda Journal of Agriculture Science*, 13: 25-34.
- Khampa, S., P. Chaowarat, R. Singhalert, M. Wanapat. 2009. Supplementation of yeast fermented cassava chip as a replacement concentrate on rumen fermentation efficiency and digestibility on nutrients in cattle. *Asian Journal of Animal Science*, 3: 18-24.
- Mahgoub, O., C.D. Lu, M.S. Hameed, A. Richie, A.S. Al-Halhali, K. Annamalai. 2005. Performance of Omani goats fed diets containing various metabolizable energy densities. *Small Ruminant Research*, 58: 175-180.
- Marichala, A., N. Castroa, J. Capoteb, M.J. Zamoranoa, A. Arguñelloa. 2003. Effects of live weight at slaughter (6, 10 and 25 kg) on kid carcass and meat quality. *Livestock Production Science*, 83: 247–256.
- Marques, C.A.T., A.N. de Medeiros, R.G. Costa, F.F.R. de Carvalho, M.J. de Araújo, J.N.D. Torreão. 2014. Performance and carcass traits of Moxotó growing goats supplemented on native pasture under semiarid conditions. *Revista Brasileira de Zootecnia*, 43:151-159.
- Morgan, N.K., M. Choct. 2016. Cassava: nutrient composition and nutritive value in poultry diets. *Animal Nutrition*, 2: 253-261.
- Nehring, K., G.W.F. Haelien. 1973. Feed evaluation and calculation based on net energy. *Journal of Animal Science*, 36: 949 – 963.
- Ocheja, J.O., B.C. Alex, S. Onoja, G.O. Usman, J. Ocheni, B.C. Lalabe. 2014. Performance, haematological and serum biochemical profile of weaner West African dwarf goats fed with diets containing graded levels of cashew nut shell. *International Journal of Research in Agriculture and Forestry*, 1(1): 27-33.
- Oduguwa, B.O., T.A. Amole, N. Okwelum, A.O. Oni, P.D. Toviesi, O.A. Oderinwale, O.A. Soyomokun. 2013. Performance, storability and microbiological assay of pelletized and un-pelletized cassava based diets fed to Muturu calves. *African Journal of Agriculture Research*, 8: 2750-2756.
- Oelker, E.R., C. Reveneau, J.L. Firkins. 2009. Interaction of molasses and monensin in alfalfa hay- or corn silage-based diets on rumen fermentation, total tract digestibility, and milk production by Holstein cows. *Journal of Dairy Science*, 92: 270-285.
-

- Ogundipe, S.O., P.O. Abeke, A.A. Sekon, I.I. Defwang, L.A. Adeyinka. 2003. Effects of duration of cooking on the utilization of *Lablab purpureus* beans by pullet chicks. Proceedings of the 36th Annual Conference of Nigerian Society of Production held at Ibadan, Nigeria.
- Oni, A.O., O.M. Arigbede, O.S. Sowande, U.Y. Anele, O.O., Oni, C.F.I. Onwuka, O.S. Onifade, K.O. Yusuf, P.A. Dele, R.Y. Aderinboye. 2012. Haematological and serum biochemical parameters of West African Dwarf goats fed dried cassava leaves-based concentrate diets. *Tropical Animal Health and Production*, 44: 483–490.
- Oni, A.O., A. Abatan, K. Adebayo, O.S. Sowande, S. Iposu, C.F.I. Onwuka 2017. Effects of supplementing cassava peels with cassava leaves and cowpea haulms on the rumen environment and blood profile parameters of West African dwarf goats. *Archives of Zootechnia*, 66: 395-402.
- Onyeonagu, C.C., S.M. Eze. 2013. Proximate compositions of some forage grasses and legumes as influenced by season of harvest. *African Journal of Agriculture Research*, 8: 4033-4037.
- Rodriguez, A.A., E.E. González, P.F. Randel. 2014. Performance and meat carcass characteristics of locally slaughtered sheep and goats raised by grazing native tropical grasses with or without supplementation. *Journal of Agricultural University P.R.*, 98:129-146.

Received: 14 September 2018

Accepted: 17 November 2018